

What is claimed is:

1. A dosage form for the oral administration of a methylphenidate drug, comprising two groups of particles, each containing said drug, wherein:
 - a) said first group of particles provides a substantially immediate dose of said drug upon ingestion by a mammal, and
 - b) said second group of particles comprises coated particles, said coated particles comprising from about 2% to about 75% by weight of said drug in admixture with one or more binders, said coating comprising a pharmaceutically acceptable ammonio methacrylate in a quantity sufficient to provide a dose of said medication delayed by from about 2 hours to about 7 hours following said ingestion.
2. The dosage form of claim 1 wherein said first group of particles comprises a pharmaceutically acceptable salt of methylphenidate in powder form.
3. The dosage form of claim 1 wherein said second group of particles comprises coated particles comprising a pharmaceutically acceptable salt of methylphenidate.
4. The dosage form of claim 2 wherein the amount of said pharmaceutically acceptable salt of methylphenidate in said first group of particles is from about 2% to about 99% by weight, based on the weight of said particles.
5. The dosage form of claim 4 wherein said pharmaceutically acceptable salt of methylphenidate comprises *dl-threo* methylphenidate hydrochloride.
6. The dosage form of claim 3 wherein said pharmaceutically acceptable salt of methylphenidate comprises *dl-threo* methylphenidate hydrochloride.

08892190-071497

25

- a) particles comprising from about 2% by weight to about 99% by weight of said methylphenidate drug, in admixture with one or more binders,
- b) a coating exterior to said methylphenidate drug, comprising an ammonio methacrylate copolymer in a quantity sufficient to provide a dose of said methylphenidate delayed by from about 2 hours to about 7 hours following administration, and
- c) on the exterior surface of said coating, a layer comprising said methylphenidate drug, to provide a substantially immediate dose of said methylphenidate upon administration.

27

24. The dosage form of claim 23 wherein said methylphenidate is *dl-threo*-methylphenidate hydrochloride.

25. The dosage form of claim 23 wherein said methylphenidate is *d-threo*-methylphenidate hydrochloride.

26. The dosage form of claim 23 wherein said coating comprises a first ammonio methacrylate copolymer comprising, as polymerized units, acrylic groups and trimethylammonioethyl methacrylate in a ratio of from about 30:1 to about 40:1, and a second ammonio methacrylate copolymer comprising, as polymerized units, acrylic groups and trimethylammonioethyl methacrylate in a ratio of from about 15:1 to about 20:1.

27. A dosage form for the oral administration of *d-threo*-methylphenidate hydrochloride comprising two groups of particles, each containing *d-threo*-methylphenidate, wherein:

- a) said first group of particles comprises *d-threo*-methylphenidate hydrochloride and provides a substantially immediate dose of said *d-threo* methylphenidate upon ingestion by a mammal, and
- b) said second group of particles comprises coated particles, said coated particles comprising from about 2% to about 75% by weight of *d-threo*-methylphenidate hydrochloride in admixture with one or more binders, said coating comprising a pharmaceutically acceptable ammonio methacrylate copolymer in an amount sufficient to provide a dose of said *d-threo*-methylphenidate delayed by from about 2 hours to about 7 hours following said ingestion.

28. A method for treating disease in a patient in need of treatment comprising administering to the patient a dosage form providing once-daily oral administration of

08992190 074497

28

d-threo-methylphenidate hydrochloride, said dosage form comprising two groups of particles, each containing *d-threo*-methylphenidate, wherein:

- a) said first group of particles comprises from about 2% to about 99% by weight of *d-threo*-methylphenidate hydrochloride and provides a substantially immediate dose of said *d-threo* methylphenidate upon ingestion by a mammal; and
- b) said second group of particles comprises coated particles, said coated particles comprising from about 2% to about 75% by weight of *d-threo*-methylphenidate in admixture with one or more binders, and a coating consisting of an ammonio methacrylate copolymer in an amount sufficient to provide a dose of said *d-threo*-methylphenidate hydrochloride delayed by from about 4 hours to about 7 hours following said ingestion.

28
29. A dosage form of a pharmaceutically acceptable salt of *d-threo*-methylphenidate providing an *in vitro* release profile comprising two pulses of drug release, wherein said pulses are temporally separated by from about 2 hours to about 7 hours.

29
30. A dosage form of a pharmaceutically acceptable salt of *d-threo*-methylphenidate providing an *in vivo* plasma concentration of said *d-threo*-methylphenidate comprising two maxima, wherein said maxima are temporally separated by from about 2 hours to about 7 hours, and wherein the magnitude of said maxima differ by no more than about 30 percent.

30
31. A dosage form according to claim 23 wherein said ammonio methacrylate copolymer comprises a first copolymer of methyl methacrylate, ethyl acrylate and TAMCl in a ratio of 2:1:0.1 and a second copolymer of methyl methacrylate, ethyl acrylate, and TAMCl in a ratio of 2:1:0.2.

29

08992190.071497